



Cepetor™

DIN 02337177

Medetomidine* hydrochloride injection - 1 mg/mL
Sedative and analgesic for dogs

Cepetor™ is indicated for use as a sedative and analgesic to facilitate clinical examinations, clinical procedures, minor surgical procedures with the exception of those requiring muscle relaxation, and minor dental procedures where intubation is not required, in dogs over 12 weeks of age. The IV route of administration is more efficacious for dental care.

Dosage and Administration:

Cepetor™ should be administered at the rate of 750 µg IV or 1,000 µg IM per square meter of body surface. See chart to the right.

See package insert for full information on side effects, precautions, warnings and contraindications.

Storage: 15-30° C. Protect from freezing.

*Each mg of medetomidine contains 0.5mg of dexmedetomidine (active ingredient) and 0.5mg of levomedetomidine (inactive compound)

Product Codes:	CDMV	VPCL	WDDC	AVP	MVP
Cepetor 10mL -	109025	7750900	120827	3820470	7220025
Revertor 10mL -	109026	7754600	120828	3823462	7610015

Revertor™

DIN 02337207

Atipamezole hydrochloride injection - 5 mg/mL
Medetomidine reversal agent for dogs

Revertor™ is indicated for the reversal of the clinical effects of the sedative and analgesic agent, medetomidine hydrochloride, in dogs.

Dosage and Administration:

Revertor™ is administered intramuscularly regardless of the route used for medetomidine hydrochloride. The concentration of Revertor™ has been formulated such that the volume of injection is the same (mL for mL) as the recommended dose volume of medetomidine hydrochloride, and may be given at any time following medetomidine hydrochloride administration. Although injection volumes are the same, the concentration of Revertor™ (5.0 mg/mL) is 5 times that of medetomidine hydrochloride (1.0 mg/mL). Dogs that are sedated but ambulatory may be treated with Revertor™, if warranted. The dosage of Revertor™ is calculated based upon body surface area. See chart to the right.

See package insert for full information on side effects, precautions, warnings and contraindications.

Storage: 15-30° C. Protect from light.

Body weight (kg) IV Administration	Injection Volume (mL)	Body weight (kg) IM Administration
1.5-2.2	0.1	1.8-2.3
2.3-3.5	0.15	2.4-3.3
3.6-5.1	0.2	3.4-4.5
5.2-6.9	0.25	4.6-6.4
7.0-9.9	0.3	6.5-9.4
10.0-14.4	0.4	9.5-12.7
14.5-19.5	0.5	12.8-16.3
19.6-25.1	0.6	16.4-20.2
25.2-31.1	0.7	20.3-24.4
31.2-37.6	0.8	24.5-28.9
37.7-44.4	0.9	29.0-36.1
44.5-55.3	1.0	36.2-46.3
55.4-71.1	1.2	46.4-57.3
71.2-88.2	1.4	57.4-75.8
88.3+	1.5	75.9+
	2	



Body weight (kg) if medetomidine is given IV	Revertor injection Volume (mL) IM	Body weight (kg) if medetomidine is given IM
1.5-2.2	0.1	1.8-2.3
2.3-3.5	0.15	2.4-3.3
3.6-5.1	0.2	3.4-4.5
5.2-6.9	0.25	4.6-6.4
7.0-9.9	0.3	6.5-9.4
10.0-14.4	0.4	9.5-12.7
14.5-19.5	0.5	12.8-16.3
19.6-25.1	0.6	16.4-20.2
25.2-31.1	0.7	20.3-24.4
31.2-37.6	0.8	24.5-28.9
37.7-44.4	0.9	29.0-36.1
44.5-55.3	1.0	36.2-46.3
55.4-71.1	1.2	46.4-57.3
71.2-88.2	1.4	57.4-75.8
88.3+	1.5	75.9+
	2	