



Cepesedan™

Detomidine hydrochloride injection - 10 mg/mL

Sedative and analgesic for horses and yearlings

Cepesedan™ is indicated for use as a sedative and analgesic to facilitate minor surgical and diagnostic procedures in mature horses and yearlings.

Dosage and Administration:

Administer Cepesedan™ intravenously or intramuscularly at the rates of 20 or 40 µg detomidine hydrochloride per kg of body weight (0.2 to 0.4 mL of Cepesedan™ per 100 kg or 220 lbs.), depending upon the depth and duration of sedation and analgesia required. Analgesia is more pronounced when the drug is given intravenously. Sedative and analgesic effects will usually occur in 2 to 5 minutes and will persist for 30 minutes to 2 hours, depending upon the dosage and route of administration.

Prior to and following injection, the animal should be allowed to rest quietly. The full effect should be reached within 2 to 4 minutes after I.V. injection and 3 to 5 minutes after intramuscular injection.

See package insert for full information on side effects, precautions, warnings and contraindications.

Packaging:

5 and 20 mL multidose vials

Storage:

15° - 30° C. Protect from light.

Orders & Product Information:

Call 1 888 590 9839

Product Codes:

	CDMV	VPCL	WDDC	AVP	MVP
Cepesedan 5mL-	109023	7750800	120825	3820460	9640005
Cepesedan 20mL-	109024	7750802	120826	3820462	9640006



Another product brought to you by Modern Veterinary Therapeutics, your horse authority.