



Nerfasin 20™



Xylazine injection - 20 mg/mL

DIN 02444089

Nerfasin 20™ should be used in dogs, cats, and cattle when it is desirable to produce a state of sedation accompanied by a shorter period of analgesia.

Dosage and Administration: Dogs & Cats

Intravenously - 0.5 mL/20 lbs of body weight (0.5 mg/lb) (1.1 mg/kg)
Intramuscularly or subcutaneously - 1.0 mL/20 lbs body weight (1 mg/lb or 2.2 mg/kg). In large dogs (over 50lbs.) a dosage of 0.5 mg/lb administered intramuscularly may provide sufficient sedation and/or analgesia for most procedures.

Dosage and Administration: Cattle

Intramuscularly - Range of 0.25 to 0.75 mL/100 lbs body weight (Equivalent to 0.05 to 0.15 mg/lb or 0.11 to 0.33 mg/kg).

See package insert for full information on side effects, precautions, warnings and contraindications.

Storage: Store at room temperature, 15-30°C; Do not freeze. Store the vial in the upright position. Discard any remainder 28 days after broaching. The stopper should not be punctured more than 20 times after broaching

Packaging: 25 mL multidose vial

Orders & Product Information:
Call 1 888 590 9839

	CDMV	VPCL	WDDC	AVP	MVP
25mL:	118550	7753260	131618	3822640	9690008

